

Yellow Bristle Grass

• Control in Pasture

Panther[®] S. A selective grass herbicide for the control of yellow bristle grass in pasture.

User Instructions

Use 750 mL/ha of Panther S. If applied by;

- Boom sprayer, apply in 250L/ha of water.
- Handgun, apply at 75 mL per 100L, using 1000L/ha of water. Uniform coverage is required.
- Knapsack, apply at 10 mL per 10L, using 750L/ha of water. Uniform coverage is required.

Do not use with an adjuvant or any other agrichemical.
Do not use Panther S within 5 days of any other herbicide treatment.

Apply during December/January.

Apply 7 days after grazing.

Avoid spray overlap or over application.

Apply to established pasture only.

The withholding period from application to grazing is 28 days.

This product is to be used for yellow bristle grass control only. This product does not control broadleaf weeds. For broadleaf weed control, a separate spray programme is required.



Collect your AgRewards points from this product. Look out for the AgRewards stickers on packs.
Visit www.agrewards.co.nz to find out more.



Panther[®] S. A selective grass herbicide for the control of yellow bristle grass in pasture.

GENERAL INFORMATION

Panther S is a selective grass herbicide for use on spring or autumn sown wheat and perennial ryegrass for the control of wild oats and lesser canary grass (*Phalaris minor*), and for use in pasture for the control of yellow bristle grass.

DIRECTIONS FOR USE

Panther S can be safely used on crops undersown with clover and with certain grasses (see 'Weed Control Notes' below). Do NOT use on crops undersown with Concord ryegrass.

CROP	RATE	WEED	CRITICAL COMMENTS
Wheat	750 mL/ha in 250 L/ha of water. Use with Synoil at 0.5% of the spray mix (500 mL per 100 L)	Wild Oats Lesser Canary Grass (<i>Phalaris minor</i>)	Apply when wild oats are actively growing, and are at a growth stage between 3 leaf and 1st node visible. Apply when lesser canary grass are actively growing, and are at a growth stage between 3 leaf and the end of tillering. Avoid application between periods of moisture stress.
Perennial Ryegrass (<i>Lolium perenne</i>)	750 mL/ha in 250 L/ha of water. Do NOT use with an adjuvant or any other agrichemical	Wild Oats	Apply in early spring when wild oats are actively growing, and are at a growth stage between 3 leaf and the end of tillering. NOTE: Delaying treatment is more likely to result in lower seed yields.
Pasture (Established)	750 mL/ha in 250 L/ha of water. Do NOT use with an adjuvant or any other agrichemical	Yellow Bristle Grass	Apply during December/January. Apply 7 days after grazing.

WEED CONTROL NOTES: The following weeds are not controlled by Panther S: Annual and hybrid ryegrasses, browntop, chewings fescue, cocksfoot, gnawed canary grass (*Phalaris paradoxa*), grazing brome (*Bromus stamineus*), prairie grass, tall fescue, and most broadleaf weeds.

(For complete instructions, refer to the product label).

www.orionagriscience.co.nz

For more information contact Orion AgriScience Ltd. Freephone: 0800 674 6627

Panther[®] is a Registered Trade Mark of Orion AgriScience Ltd, Registered pursuant to the ACVM Act 1997, No. P8064. See www.foodsafety.govt.nz for registration conditions. See www.epa.govt.nz for approval conditions.

