



orion AGRISCIENCE



HARMFUL | ECOTOXIC | CORROSIVE
KEEP OUT OF REACH OF CHILDREN

AlignTM

**PLANT GROWTH
REGULATOR**

A growth regulator used to shorten and stiffen the straw of cereal crops to improve the resistance to lodging.

Active Ingredient:

305 g/litre mepiquat chloride and 155 g/litre chlorthalipron in the form of a soluble concentrate.



9 421004 531030

20 Litres





PLEASE READ THE LABEL COMPLETELY BEFORE USE

HAZARD CLASSIFICATIONS: 6.1D, 6.9B, 8.1A, 9.1C, 9.2B, 9.3B

WARNING: HARMFUL, CORROSIVE, ECOTOXIC

May be harmful if swallowed, inhaled or absorbed through the skin. May cause organ damage following repeated ingestion at high doses. Corrosive to aluminium. Harmful to aquatic organisms. Toxic to soil organisms and terrestrial vertebrates.

PRECAUTIONS

STORAGE: Keep out of reach of children. Store in the original, tightly closed container, in a secure place away from foodstuffs. Storage of certain quantities of this product trigger specific storage requirements (Refer to Safety Note). When stored appropriately this product should show no significant degradation for two years from the date of manufacture. Contact your supplier for further information about the use of product that is older than this.

PERSONAL PROTECTION: Avoid contact with eyes and skin. Avoid inhalation of spray mist. When mixing or applying, wear overalls, boots, gloves, and eye protection. Clean protective clothing daily after work. Do not eat, drink or smoke while using. Remove protective clothing and wash hands and face thoroughly before meals and after work.

USE: Apply with well maintained and calibrated spray equipment.

DISPOSAL: Dispose of this product only by using in accordance with this label, or at a suitable landfill. Triple rinse container and add rinsate to the spray tank. Dispose of container to a suitable landfill or agricultural recycling scheme. Do not burn. Do not use container for any other purpose.

ENVIRONMENTAL: Use this product carefully. Do not contaminate aquatic environments with product, spray drift or empty container. Do not allow spray drift outside the target area to occur.

First Aid

If swallowed do not induce vomiting. If splashed in eyes, wash out immediately with running water for several minutes. If skin or hair contact occurs, remove contaminated clothing and wash with soap and running water. If inhaled, immediately move to fresh air and rest. Maintain half upright position if breathing is difficult. Call the National Poisons Centre **0800 POISON (0800 764 766)** (24 hrs) or a doctor immediately for advice.

Emergency Response

In a transport emergency call **111, Police or Fire Brigade**. For specialist advice in an emergency only, call **0800 CHEMCALL (0800 243 6225)** (24 hrs).

Spillage

Exclude non-essential people from the area. Wear protective clothing and prevent material from entering waterways. Absorb spill with inert material such as soil, sand, etc, and place in a waste container. Wash area with water and detergent, and absorb with further inert material. Dispose of safely to a suitable landfill.

Safety Data Sheet

Additional information is provided in the safety data sheet.

Freephone **0800 ORIONAS (0800 6746627)** or visit www.orionagriscience.co.nz.

GENERAL INFORMATION

Align reduces the risk of lodging in vigorous crops, susceptible cultivars, high nitrogen use situations and in field situations where there is a history of lodging. Early treatment provides greater protection from stem lodging, whereas later applications reduce yield losses caused by stem and neck breaking.

Ingredients: Align contains 305 g/litre mepiquat chloride and 155 g/litre chlorthethphon as the active ingredients, and up to 14g/L of acetic acid.

CAUTION:

To work effectively, crops to be treated should be actively growing and free of stress caused by nutrient or moisture deficiency. Do NOT apply when temperatures are above 21°C, or to crops suffering damage or physical stress from frost, hail or herbicide damage. Secondary tillering may be a problem in treated crops, especially in situations where the application has been made early followed by a period of stress. This may have implications, especially in crops grown for malting barley, or in other situations where a bold sample of grain is required.

APPLICATION METHODS

Obtain uniform spray coverage using conventional ground boom sprayer and aerial equipment. Flush equipment thoroughly with clean water after use.

DIRECTIONS FOR USE

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds. It is an offence to use this product on animals.

Crop	Rates per Hectare	Critical Comments
Autumn Sown Barley, Ryecorn and Triticale	1.0 - 1.5 litres in 200 litres of water. Always add 25ml/100 litres water of Synergy™ Wetter Sticker.	Apply from Zadoks GS 37 (flag leaf visible) through until GS 49 (first awns visible). Use the higher rate where the risk of lodging or straw and neck breaking is high.
Spring Sown Barley	1.0 - 1.5 litres in 200 litres water. Always add 25ml/100 litres water of Synergy Wetter Sticker.	Apply to stress free crops from Zadoks GS 39 (flag leaf ligule visible) through until GS 49 (first awns visible). Use the higher rate where the risk of lodging or straw and neck break is high. Use the low rate in malting barley, only in situations when the crop is free from stress.

MIXING

When used in a mixture always add Align to the spray tank last. Use spray solution immediately. Where water has a high pH, or water containing calcium salts cannot be avoided, the addition of a pH Buffer is recommended to reduce chemical deactivation and hydrolysis.

COMPATIBILITY

May be tank mixed with most commonly used insecticides and fungicides. Do not mix with herbicides.

RAINFALL

Do not apply if rain is expected within 6 hours of spraying, or if the crop is wet, or frost is expected.

WARRANTY

This formulation is warranted to contain the specified active ingredient within accepted analytical tolerance when packed. The user bears the risk for damage resulting from factors beyond the manufacturer's control. The manufacturer declines all liability for damage resulting from improper storage or use of the product.

Registered pursuant to the ACVM Act 1997, No. P5989

See www.foodsafety.govt.nz for registration conditions.

Approved pursuant to the HSN0 Act 1996, No. HSR000806

See www.epa.govt.nz for approval conditions.

™ ALIGN and SYNERGY are Trade Marks of Orion AgriScience Ltd.

Registrant: Orion AgriScience Ltd,

Unit 1, 15 Sir Gil Simpson Drive,

Harewood, Christchurch 8053

www.orionagriscience.co.nz.



UN 3265

CORROSIVE LIQUID, ACIDIC, ORGANIC, N.O.S. (CHLORETHEPHON 15%) PACKING GROUP III

HAZCHEM CODE 2X

ADDITIONAL INFORMATION: CORROSIVE TO ALUMINIUM

IN A TRANSPORT EMERGENCY, DIAL 111 POLICE OR FIRE BRIGADE.

FOR SPECIALIST ADVICE IN AN EMERGENCY CALL 0800 CHEMCALL (0800 243 622)

orion
AGRISCIENCE

ALIGN Batch No: Date of manufacture:

