

orion AGRISCIENCE

WARNING

KEEP OUT OF REACH OF CHILDREN



TriumphTM
2G

GRANULAR HERBICIDE



For control of blackberry, broom, gorse,
ragwort, sweet brier, thistles, tutsan and
other difficult to kill perennial weeds

Active Ingredient:

20 g/kg picloram as the amine salt in the form of a pellet



9 421004 532167

5 Kg

140 x 140mm (WxH)

PLEASE READ THE LABEL COMPLETELY BEFORE USE**HAZARD CLASSIFICATIONS:****Skin sensitisation Category 1, Hazardous to soil organisms, Hazardous to the aquatic environment chronic Category 3****WARNING: ECOTOXIC**

May cause an allergic skin reaction. Harmful to aquatic life with long lasting effects. Very toxic to the soil environment.

PRECAUTIONS**HANDLING REQUIREMENTS:**

Records of use as described in NZS 8409 Management of Agrichemicals must be kept if 3 kg or more is applied within 24 hours and likely to leave the application area.

PERSONAL PROTECTION:

Avoid contact with eyes and skin. Avoid inhalation of dust. When applying, wear overalls, boots, gloves and eye protection. Clean protective clothing daily after work. Do not eat, drink or smoke while using. Remove protective clothing and wash hands and face thoroughly before meals and after work.

STORAGE:

Keep out of reach of children. Store in the original, tightly closed container, in a secure place. When stored appropriately this product should show no significant degradation for two years from the date of manufacture. Contact your supplier for further information about the use of product that is older than this.

DISPOSAL:

Dispose of this product only by using in accordance with this label, or at a suitable landfill. Dispose of container to a suitable landfill. Do not burn. Do not use container for any other purpose.

ENVIRONMENTAL:

Use this product carefully. Take all reasonable steps to ensure that this product does not cause any significant adverse effects to the environment beyond the application area. Do not apply directly into or onto water. Do not contaminate aquatic environments with product or empty container. Collect spillage.

EMERGENCY RESPONSE

In a transport emergency call **111, Police or Fire Brigade**. For specialist advice in an emergency only, call **0800 CHEMCALL (0800 243 6225)** (24 hrs).

SPILLAGE

Exclude non-essential people from the area. Wear protective clothing and prevent material from entering waterways. Moisten to prevent dusting and sweep up. Place in a waste container. Dispose of safely to a suitable landfill.

SAFETY DATA SHEET

Additional information is provided in the safety data sheet. Freephone **0800 ORIONAS (0800 674 6627)** or visit **www.orionagriscience.co.nz**.

WARRANTY

This formulation is warranted to contain the specified active ingredients within accepted analytical tolerance when packed. The user bears the risk for damage resulting from factors beyond the manufacturer's control. The manufacturer declines all liability for damage resulting from improper storage or use of the product.

Registered pursuant to the ACVM Act 1997, No. P7936
See www.foodsafety.govt.nz for registration conditions.
Approved pursuant to the HSN0 Act 1996, No. HSR000554
See www.epa.govt.nz for approval controls.

TMTRIUMPH is a Trade Mark of Orion AgriScience Ltd

FIRST AID

If medical advice is needed, have the product container or label at hand. If swallowed do not induce vomiting. If splashed in eyes, wash out immediately with running water for several minutes. If skin or hair contact occurs, remove contaminated clothing and wash with soap and running water. If inhaled, immediately move to fresh air and rest. Maintain half upright position if breathing is difficult. Call the National Poisons Centre **0800 POISON (0800 764 766)** (24 hrs) or a doctor immediately for advice.

Batch No:

Date of manufacture:

TRIUMPH 2G

The logo for Orion AgriScience, featuring the word "orion" in a stylized, lowercase font with a swoosh above it, and "AGRISCIENCE" in a smaller, uppercase font below it. The logo is set against a dark background with a white border.

0325

PLEASE READ THE LABEL COMPLETELY BEFORE USE

GENERAL INFORMATION

Triumph 2G is a herbicide for spot application onto brushweeds and broadleaf weeds in areas where spraying is impractical or undesirable.

Note: Triumph 2G is damaging to clovers. The area to be treated should be kept to a minimum. **Do not use on peat or heavy clay soils.**

DIRECTIONS FOR USE

RATES OF APPLICATION AND TIMING

Weed	Rate	Critical Comments
Docks, nodding thistle, ragwort	2 g per plant	Apply half a level teaspoon to the crushed centre of each plant
Broom, fennel, goats rue, hemlock, inkweed, lupin, Mexican devil weed, St Johns wort, woolly nightshade	30 g per sq metre	Best results are achieved on light sandy soils or stony soils, with application being made in late winter to late spring
Barberry (small bushes), blackberry, boxthorn (seedling), gorse, heath, Himalayan honeysuckle, horehound, matagouri, sweet brier, tutsan, tutu	55 g per sq metre	

Best results are obtained under favourable growth conditions. Sufficient soil moisture must be present at the time of application to move the active ingredient into the soil. Application under dry conditions will give poor results.

APPLICATION METHODS

Triumph 2G should be applied over the top of the plant, and in the case of brushweeds, extending out to 30 cm beyond the dripline. On steep slopes, more granules should be applied on the uphill side as some downward leaching will occur.

WITHHOLDING PERIODS

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

Poisonous plants may become more palatable after treatment, and livestock should be kept out until plants have died down. Otherwise, no withholding period is required.

RESISTANCE MANAGEMENT

GROUP	4	HERBICIDE
-------	---	-----------

Triumph 2G contains the active ingredient picloram, which belongs to the pyridine carboxylate chemical family. Some naturally occurring weeds may become resistant when excessive use of the same herbicide occurs. To prevent or delay the development of resistance, use in tank mixes and/or rotate use with herbicides with a different mode of action. Consult <http://resistance.nzpps.org/index.php> for alternative modes of action and details of resistance management strategies (if any) for the uses listed on the label.