

The logo features the word "orion" in a large, white, lowercase sans-serif font, followed by "AGRISCIENCE" in a smaller, white, uppercase sans-serif font. The text is set against a green background that has a white swoosh element on the left side.

orion

AGRISCIENCE

DANGER

KEEP OUT OF REACH OF CHILDREN



LegendTM

HERBICIDE

For the control of broadleaf weeds in WHEAT, BARLEY, OATS and RYEGRASS SEED CROPS as listed in the Situations for Use Table

Active Ingredient:

600 g/litre mecoprop, 150 g/litre MCPA and 18.7 g/litre dicamba in the form of a soluble concentrate



9 421004 531641

20 Litres

READ THE LABEL BEFORE USE

HAZARD CLASSIFICATIONS:

Acute Oral Toxicity (Cat. 4), Skin Sensitisation (Cat 1), Eye Damage (Cat. 1), Specific Target Organ Toxicity Repeat Exposure (Cat. 1), Hazardous to the Aquatic Environment Acute (Cat. 1), Hazardous to the Aquatic Environment Chronic (Cat. 1), Hazardous to Soil Organisms, Hazardous to Terrestrial Vertebrates

DANGER: TOXIC, ECOTOXIC

Harmful if swallowed. May cause an allergic skin reaction. Causes serious eye damage. May cause damage to organs through prolonged or repeated exposure. Very toxic to aquatic life with long lasting effects. Hazardous to soil organisms (selective herbicide). Hazardous to terrestrial vertebrates.

PRECAUTIONS

HANDLING REQUIREMENTS:

This product must only be handled, mixed and applied by a suitably qualified person as defined under the HPC notice 2017 Part 4C. Records of use as described in NZS 8409 Management of Agrichemicals must be kept if 3 L or more is applied within 24 hours and likely to leave the application area.

PERSONAL PROTECTION

Avoid contact with eyes and skin. Do not breathe in the spray mist. When mixing, loading or applying, wear overalls, boots, gloves and eye protection. Wear respiratory protection if spray mist could be inhaled. Take off contaminated clothing and wash it before reuse. Contaminated work clothing should not be allowed out of the workplace. Do not eat, drink or smoke when using this product. Wash hands thoroughly after handling.

STORAGE

Store in the original container. Keep container tightly closed. When stored appropriately this product should show no significant degradation for two years from the date of manufacture. Contact your supplier for further information about the use of product that is older than this.

DISPOSAL

Dispose of this product by using it in accordance with this label. Dispose of the container via AgRecovery after triple rinsing (use the rinsate for spraying). Do not burn. Do not use the container for any other purpose.

ENVIRONMENTAL

Take all reasonable steps to ensure that this product does not cause any significant adverse effects to the environment beyond the application area. Do not apply directly into or onto water. Collect spillage. Spray drift or equipment contamination may cause serious damage to desirable plants and susceptible crops such as vegetables, flowers, fruit trees, ornamental shrubs, grapes and tomatoes. Do not use the same equipment on other susceptible crops unless decontaminated completely.

FIRST AID

If medical advice is needed, have product container or label at hand.

IF SWALLOWED: Rinse mouth. Get medical help.

IF ON SKIN: Wash with plenty of water. If skin irritation or rash occurs: Get medical help.

IF IN EYES: Immediately rinse with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing. Get medical help.

Get medical help if you feel unwell. For advice, call the Poisons Centre on **0800 POISON (0800 764 766)** or a Doctor.

EMERGENCY RESPONSE

In an emergency call **111, Police or Fire Brigade**. For specialist advice in an emergency only, call **0800 CHEMCALL (0800 243 6225)** (24 hrs).

SAFETY DATA SHEET

Additional information is provided in the safety data sheet. Freephone **0800 ORIONAS (0800 674 6627)** or visit **www.orionagriscience.co.nz**

DIRECTIONS FOR USE

Apply with well-maintained and calibrated spray equipment. Application must be by a qualified loader, contractor or person.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds

Apply when the cereal crop is at the 4 leaf stage. Apply Legend from late autumn to early winter in ryegrass seed crops when the grass is at least at the 4 - 6 leaf stage of growth. Do not apply after the cereal and ryegrass seed crops have reached the 'jointing' or 'node' stage. Best results are obtained when the weeds are actively growing and not under stress. For consistent results, water volumes should not be less than 150 litres per hectare. Do not exceed the recommended application rate. Do not overlap or "spray out" corners when spraying. Do not apply to crops undersown with clovers or to any cereal crop from which the straw may be required for horticultural mulching.

Batch No:

Date of manufacture:

SITUATIONS FOR USE

WEED TABLE	CROP	RATE PER HECTARE	CRITICAL COMMENTS
Annual nettle, black nightshade, mouse eared chickweed, catchfly, cornbind, docks, fathen, hedge mustard, mallow spp, redroot, spurrey, shepherd's purse, thistles (noddling, scotch, sow, winged and californian thistle regrowth from rootchips), twincross, wireweed	Wheat, Barley, Oats, Ryegrass Seed Crops	3 – 3.5 litres per hectare using 150 - 200 litres of water.	Apply to seedling weeds. Use the higher rate when weeds are at the 6 to 8 true leaf stage.
Chamomiles, cleavers, fumitory, scrambling speedwell, stinking mayweed, water pepper, yarrow		3.5 – 4 litres per hectare using 150- 200 litres of water.	Apply to seedling weeds. Use the higher rate when weeds are at or over the 6 true leaf stage. Use 4 litres per hectare for chamomiles, stinking mayweed, water pepper and yarrow.

MIXING

Add the required quantity of Legend to spray tank partly filled with water and maintain agitation throughout filling and spraying. The use of additional wetting agent is not recommended.

COMPATIBILITY

Legend is compatible with most commonly used herbicides.

RAINFALL

Do not spray if rain is expected within 6 hours of spraying, however 12 hours drying time is preferable.

RESISTANCE MANAGEMENT

GROUP	4	HERBICIDE
-------	---	-----------

Legend contains mecoprop and MCPA, belonging to the phenoxy chemical family, and dicamba, belonging to the benzoic acid chemical family. Some naturally occurring weeds may become resistant when excessive use of the same herbicide occurs. To prevent or delay the development of resistance, use in tank mixes and/or rotate use with herbicides with a different mode of action. Consult <http://resistance.nzpps.org/index.php> for alternative modes of action and details of resistance management strategies (if any) for the crops listed on the label.

WARRANTY

This formulation is warranted to contain the specified active ingredient within accepted analytical tolerance when packed. The user bears the risk for damage resulting from factors beyond the manufacturer's control. The manufacturer declines all liability for damage resulting from improper storage or use of the product.

Registered pursuant to the ACVM Act 1997, No. P5005
See www.foodsafety.govt.nz for registration conditions.
Approved pursuant to the HSNO Act 1996, No. HSR000353
See www.epa.govt.nz for approval controls.

™Legend is a Trade Mark of Orion Agriscience Limited.

Registrant: Agsin Pte Ltd, Zagro Global Hub, 5 Woodlands Terrace, Singapore 738430

Distributor: Orion AgriScience Ltd, Unit 1, 15 Sir Gil Simpson Drive, Harewood, Christchurch 8053

0800 674 6627
www.orionagriscience.co.nz

LEGEND



0425

AGRISCIENCE