

100 x 110mm (WxH)

orion AGRISCIENCE



WARNING
KEEP OUT OF REACH OF CHILDREN



ChordTM

HERBICIDE

A selective herbicide for the control of buttercups and docks in PASTURE, WHEAT, BARLEY and OATS, and in CONSERVATION TILLAGE PROGRAMS

Active Ingredient:

750 g/kg of thifensulfuron-methyl in the form of a water dispersible granule



500 Gram

100 x 110mm (WxH)

PLEASE READ THE LABEL COMPLETELY BEFORE USE

PEEL HERE

HAZARD CLASSIFICATIONS:**Eye irritation Category 2, Hazardous to soil organisms, Hazardous to the aquatic environment acute Category 1, Hazardous to the aquatic environment chronic Category 1****WARNING: ECOTOXIC**

Causes serious eye irritation. Very toxic to aquatic life with long lasting effects. Very toxic to the soil environment.

PRECAUTIONS**HANDLING REQUIREMENT**

This product must only be handled, mixed and applied by a suitably qualified person as defined under the HPC notice 2017 Part 4C. Records of use as described in NZS 8409 Management of Agrichemicals must be kept if 3 kg or more is applied within 24 hours and likely to leave the application area.

PERSONAL PROTECTION:

Avoid contact with eyes and skin. Avoid inhalation of dust or spray mist. When mixing or applying, wear overalls, boots, gloves and eye protection. Do not eat, drink or smoke while using. Remove protective clothing and wash hands and face thoroughly before meals and after work.

STORAGE:

Keep out of reach of children. Store in the original, tightly closed container, in a secure place. Storage of certain quantities of this product trigger specific storage requirements (Refer to Safety Note). When stored appropriately this product should show no significant degradation for two years from the date of manufacture. Contact your supplier for further information about the use of product that is older than this.

DISPOSAL:

Dispose of this product only by using in accordance with this label, or at a suitable landfill. Dispose of packaging to a suitable landfill or agricultural recycling scheme. Do not burn. Do not use packaging for any other purpose.

ENVIRONMENTAL:

Use this product carefully. Take all reasonable steps to ensure that this product does not cause any significant adverse effects to the environment beyond the application area. Do not apply directly into or onto water. Do not contaminate aquatic environments with product, spray drift or packaging. Spray drift or equipment contamination may cause serious damage to desirable plants. Do not use or flush equipment near to desirable plants or their roots. Do not use the same equipment to apply other chemicals to susceptible crops unless thoroughly cleaned and decontaminated. Collect spillage.

FIRST AIDIf medical advice is needed, have the product container or label at hand. If swallowed do NOT induce vomiting. If splashed in eyes, wash out immediately with running water for several minutes. If skin or hair contact occurs, remove contaminated clothing and wash with soap and running water. For advice, call the National Poisons Centre **0800 POISON (0800 764 766)** (24 hrs) or a doctor immediately.**EMERGENCY RESPONSE**In a transport emergency call **111, Police or Fire Brigade**. For specialist advice in an emergency only, call **0800 CHEMCALL (0800 243 6225)** (24 hrs).**SPILLAGE**

Exclude non-essential people from the area. Wear protective clothing and prevent material from entering waterways. If granules have spilt, moisten to prevent dusting and sweep up. If liquid, absorb spill with inert material such as soil, sand, etc. Place in waste container. Wash area with water, and absorb with further inert material. Dispose of safely to a suitable landfill.

Batch No:

Date of manufacture:

SAFETY DATA SHEETAdditional information is provided in the safety data sheet. Freephone **0800 ORIONAS (0800 674 6627)** or visit www.orionagriscience.co.nz.

Registered pursuant to the ACVM Act 1997, No. P7829

See www.foodsafety.govt.nz for registration conditions.

Approved pursuant to the HSN0 Act 1996, No. HSR000236

See www.epa.govt.nz for approval controls.

TM CHORD and SILMAXX are Trade Marks of Orion Agriscience Limited

SYNERGY is a Registered Trademark of Orion Agriscience Limited

Registrant: Agsin Pte Ltd, Zagro Global Hub, 5 Woodlands Terrace, Singapore 738430**Distributor:** Orion AgriScience Ltd, Unit 1, 15 Sir Gil Simpson Drive, Harewood, Christchurch 8053.**0800 674 6627**www.orionagriscience.co.nz**CHORD**

PLEASE READ THE LABEL COMPLETELY BEFORE USE

GENERAL INFORMATION

Chord is a selective herbicide for the control of buttercups and docks in pasture and cereals. It can also be used in conjunction with glyphosate for weed control in conservation tillage programmes.

Chord works best in spring when weeds are small and actively growing. Susceptible weeds will stop growing soon after treatment, but damage symptoms may take 2 – 3 weeks to appear.

APPLICATION METHODS

Chord can be applied with conventional boom spraying equipment. Apply in a minimum of 60 litres of water per hectare. Crop injury may result from spray overlap. When starting, slowing or stopping, shut off the spray boom.

MIXING

Add the required quantity of Chord to the spray tank partly filled with water, and maintain agitation through filling and spraying. Use Chord mixtures within 24 hours of preparation as product degradation may occur.

COMPATIBILITY

Chord is compatible with most commonly used crop protection products. Note: A reduction in long term control of docks has occurred when using tank mixes with 2,4-D or MCPA.

SPRAY EQUIPMENT DECONTAMINATION

To prevent injury to subsequently sprayed sensitive crops, thoroughly decontaminate spraying equipment immediately after spraying:

1. Drain tank and flush tank, boom and hoses with clean water for at least 10 minutes.
2. Fill tank with clean water and 1 litre chlorine bleach (3% sodium hypochlorite) per 100 litres water. Keep agitation system running while flushing hoses and boom, then drain.
3. Repeat step 2.
4. Remove and clean nozzles, filters and screens separately. Rinse thoroughly with clean water and flush through hoses and boom to remove traces of chlorine.

RAINFALL

Do **NOT** spray if rainfall is expected within 4 hours of spraying.

WITHHOLDING PERIOD

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

Do **NOT** graze treated areas for 7 days after application.

WARRANTY

This formulation is warranted to contain the specified active ingredient within accepted analytical tolerance when packed. The user bears the risk for damage resulting from factors beyond the manufacturer's control. The manufacturer declines all liability for damage resulting from improper storage or use of the product.

SEE INSIDE FOR TABLE →

200 x 110mm (WxH)

PLEASE READ THE LABEL COMPLETELY BEFORE USE

RESISTANCE MANAGEMENT

GROUP	2	HERBICIDE
-------	----------	-----------

Chord belongs to the sulfonylurea chemical group. Some naturally occurring weeds may become resistant when excessive use of the same herbicide occurs. To prevent or delay the development of resistance, use in tank mixes and/or rotate use with herbicides with a different mode of action. Consult <http://resistance.nzpps.org/index.php> for alternative modes of action and details of resistance management strategies (if any) for the crops listed on the label.

DIRECTIONS FOR USE

Apply with well-maintained and calibrated spray equipment.

Do **NOT** spray pasture or cereal crops that are under stress from drought, frost, water saturated soil, disease or insect damage, as crop damage may occur. Crop damage may also occur if these conditions occur after spraying.

Do **NOT** apply to grass or clover seed crops, or hay crops, nor to crops undersown with clover or lucerne.

Crop / Situation	Application Rate	Weeds Controlled	Critical Comments
Pasture	20 g/ha (Broadcast) 1 g / 10 Litres water applied to 500 sq metres (Knapsack)	Buttercups (Annual, Creeping, Giant), Docks	Apply in spring. Apply 1 – 2 weeks after grazing so as to maximise exposure of weed seedlings. For docks, spray when in full leaf, and do NOT spray after seed heads begin to emerge or just after grazing. Follow up treatment may be necessary for well established docks and giant buttercup. NOTE: Yellowing and reduction in vigour usually occurs after application. Recovery, particularly of clovers, may take up to 4 months.
Cereals - Wheat, Barley, Oats	20 g/ha	Buttercups (Annual, Creeping), Cornbind, Docks	Apply after the crop is at the 2 leaf stage, but before the boot stage. Apply with a surfactant such as Synergy™ Wetter Sticker. Best results are obtained when applied to small actively growing weeds before canopy closure (otherwise weeds may only be suppressed).
Conservation Tillage – Cereals, Forage Brassicac, Lucerne, Maize, New Pasture, Peas, Potatoes, Squash	20 g/ha	Buttercups, Docks	Use in a tank mix with glyphosate eg. Orion Glyphosate 360, G Force® Max plus penetrant eg. Silmaxx™ or Synergy Penetrant before the crop is sown. Plantback Period: Do NOT sow the crop within 14 days of applying Chord. NOTE: Some crops may exhibit symptoms of damage, but this is transient and is unlikely to affect crop yield. The growth of perennial ryegrass may be reduced.

NOTE: Resistant weeds – Dandelion, plantains, ragwort, thistles.

SEE INSIDE FOR TABLE →