

READ THE LABEL COMPLETELY BEFORE USE**HAZARD CLASSIFICATIONS:**

Acute oral toxicity Category 4, Acute inhalation toxicity Category 4, Skin irritation Category 2, Eye irritation Category 2, Specific target organ toxicity (repeated exposure) Category 1, Hazardous to terrestrial vertebrates, Hazardous to the aquatic environment acute Category 1

DANGER: TOXIC, ECOTOXIC

Harmful if swallowed. Harmful if inhaled. Causes skin irritation. Causes serious eye irritation. Causes damage to organs through prolonged or repeated exposure. Very toxic to aquatic life. Toxic to terrestrial vertebrates.

PRECAUTIONS**HANDLING REQUIREMENTS:**

This product must only be handled, mixed and applied by a suitably qualified person as defined under the HPC notice 2017 Part 4C.

Records of use as described in NZS 8409 Management of Agrichemicals must be kept if 3 L or more is applied within 24 hours and likely to leave the application area.

PERSONAL PROTECTION

Avoid contact with eyes and skin. Do not breathe in the vapour or spray mist. When mixing or applying, wear chemical resistant overalls, boots, gloves, and eye protection. Wear a respirator if vapour or spray mist could be inhaled. Do not eat, drink or smoke during use. Clean protective equipment after use. Do not take contaminated equipment or clothing off site. Wash hands after use. Wash contaminated clothing before reuse.

STORAGE

Keep out of reach of children. Store in the original container in a cool, well ventilated and secure area. Storage of certain quantities of this product trigger specific storage requirements (Refer to Safety Note). When stored appropriately this product should show no significant degradation for two years from the date of manufacture. Contact your supplier for further information about the use of product that is older than this.

DISPOSAL

Dispose of this product by using it in accordance with this label. Dispose of the container to a suitable landfill or via AgRecovery after triple rinsing (use the rinsate for spraying). Do not burn. Do not use the container for any other purpose.

ENVIRONMENTAL

Take all reasonable steps to ensure that this product does not cause any significant adverse effects to the environment beyond the application area. Do not apply directly into or onto water. Contain and collect spillage for disposal.

FIRST AID

If medical advice is needed, have product container or label at hand.

If swallowed: Rinse mouth. Seek medical advice if feeling unwell.

If on skin: Wash with plenty of soap and water. Seek medical advice if irritation occurs.

If in eyes: Rinse cautiously with water for several minutes. Remove contact lenses, if easy to do so. Continue rinsing. Seek medical advice if irritation persists.

If inhaled: Remove to fresh air and keep at rest in a position comfortable for breathing. Seek medical advice if feeling unwell.

For advice, call the Poisons Centre on **0800 POISON (0800 764 766)** or a Doctor.

EMERGENCY RESPONSE

In a transport emergency call **111, Police or Fire Brigade**. For specialist advice in an emergency only, call **0800 CHEMCALL (0800 243 6225)** (24 hrs).

SAFETY DATA SHEET

Additional information is provided in the safety data sheet. Freephone **0800 ORIONAS (0800 674 6627)** or visit **www.orionagriscience.co.nz**

SPILLAGE

In the event of spillage, exclude non-essential people from the area. Wear protective equipment and prevent from entering waterways. Absorb spill with inert material such as soil, sand, etc, and place in a waste container. Wash area with water and detergent, and absorb with further inert material. Dispose of safely to a suitable landfill.

Batch No.

Date of Manufacture

Registered pursuant to the ACVM Act 1997, No. P9703

See www.foodsafety.govt.nz for registration conditions.

Approved pursuant to the HSNO Act 1996, No. HSR000652

See www.epa.govt.nz for approval controls.

©CALIBRE is a Registered Trade Mark of Orion AgriScience Ltd
™CONVOY is a Trade Mark of Orion AgriScience Ltd

Registrant: Agsin Pte Ltd, Zagro Global Hub,
5 Woodlands Terrace, Singapore 738430

Distributor: Orion AgriScience Ltd, Unit 1,
15 Sir Gil Simpson Drive, Harewood, Christchurch 8053.

0800 674 6627

www.orionagriscience.co.nz



CONVOY

READ THE LABEL COMPLETELY BEFORE USE

GENERAL INFORMATION

Convoy is a broad spectrum strobilurin fungicide with protectant and eradicant activity. It is locally systemic and translaminar to confer some protection to new growth. Whilst Convoy has curative activity against the early stages of infection, best results are obtained when applied prior to the onset of infection.

It is an offence to use this product on animals.

APPLICATION METHODS

Ground Application: Apply in a minimum of 150 L / ha of water.

Aerial Application: Apply in a minimum of 60 L / ha of water.

Thorough coverage of the crop is essential for best results.

MIXING

Partly fill the spray tank with water, and add the required quantity of Convoy. Maintain agitation through filling and spraying.

COMPATIBILITY

Compatible with most commonly used fungicides and insecticides.

RAINFALL

Do **NOT** apply if rain is expected within 2 hours of spraying.

WITHHOLDING PERIODS

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

Wheat, Barley (Grain)	56 days (Last application at end of ear emergence – GS 59)
Cereal (Silage/Green Feed)	28 days
Maize (Grain)	56 days
Maize (Silage/Green Feed)	28 days
Ryegrass (Seed)	35 days

READ THE LABEL COMPLETELY BEFORE USE

RESISTANCE MANAGEMENT

GROUP	11	FUNGICIDE
-------	----	-----------

Convoy contains pyraclostrobin, a fungicide from the QoI (quinone outside inhibitor) mode of action group. Resistance to this fungicide could develop from excessive use. To minimise this risk use strictly in accordance with label instructions. Apply a maximum of 2 QoI fungicides per crop per season. Always apply Convoy in a tank mix with a fungicide from a different mode of action group at the label recommended rate. Consult <http://resistance.nzpps.org/index.php> for alternative modes of action and details of resistance management strategies (if any) for the crops listed on the label.

WARRANTY

This formulation is warranted to contain the specified active ingredient within accepted analytical tolerance when packed. The user bears the risk for damage resulting from factors beyond the manufacturer's control. The manufacturer declines all liability for damage resulting from improper storage or use of the product.

DIRECTIONS FOR USE

Apply with well-maintained and calibrated spray equipment.

Crop	Disease	Rate per Hectare	Critical Comments
Barley	Leaf Rust, Net Blotch (Spot and Net form), Powdery Mildew, Ramularia Leaf and Awn Spot, Scald, Spot Blotch	800 mL plus 600 mL Calibre®	Apply as a preventative spray on susceptible varieties and when conditions favour infection. Otherwise, apply at first sign of disease. Should disease pressure persist or re-infection occur, a further application may be required 3 - 5 weeks later. Apply up to the end of ear emergence (Zadok GS 59).
Maize	Common Rust	1L	Apply as a preventative spray on susceptible varieties and when conditions favour infection. Otherwise, apply at first sign of disease. Should disease pressure persist or re-infection occur, a further application may be required 14 – 21 days later. Do NOT apply later than tasselling stage.
Ryegrass	Stem Rust	800 mL plus 600 mL Calibre®	Apply as a preventative spray on susceptible varieties and when conditions favour infection. Otherwise, apply at first sign of disease. Should disease pressure persist or re-infection occur, a further application may be required 3 - 5 weeks later. Apply up to the end of flowering (Zadok GS 69).
Wheat	Didymella Leaf Spot, Glume Blotch, Leaf Rust, Powdery Mildew, Sooty Moulds, Speckled Leaf Blotch, Stripe Rust	800 mL plus 600 mL Calibre®	Apply as a preventative spray on susceptible varieties and when conditions favour infection. Otherwise, apply at first sign of disease. Should disease pressure persist or re-infection occur, a further application may be required 3 - 5 weeks later. Apply up to the end of ear emergence (Zadok GS 59). Note: Application may be required early in the season at mid tillering for control of Speckled Leaf Blotch.