Apeggio Herbicide

NEW



Available in 1 L


Post-emergence control of certain broadleaf and grass weeds in maize and sweetcorn.

336 g/L Topramezone in the form of a suspension concentrate

Apeggio in action

Gisborne, 2025
Summer Grass and Redroot.







Target species

Broadleaf weeds (up to 8 leaves): Apple of Peru, black nightshade, broad-leaved dock (seedling), butterprint (velvetleaf), chamomile, chickweed, dandelion, fathen, fleabane, groundsel, large flowered mallow, scrambling speedwell, twin cress, white clover and willow weed.
Grass weeds (up to 3 tillers): Barnyard grass, smooth witchgrass, summer grass and yellow bristle grass.

Application

Rate & frequency	Usage guidance
200 mL/ha 1 application per year (one Litre covers 5 ha)	Apply to actively growing weeds before they reach the growth stages specified. Ensure thorough coverage for best results. Use a minimum water volume of 200 L/ha. The addition of 500 gai/ha of atrazine (e.g. 1 L/ha of a 50% liquid formulation or 555 g/ha of a 90% granular formulation) will improve the speed of knockdown and control of larger weeds. Under cool growing conditions, slight temporary chlorosis (yellowing) may occur. The crop will outgrow these symptoms, and yield will not be affected. Always use with suitable oil adjuvant unless partner product label directs otherwise.

SCAN FOR TECHNICAL SHEETS

Contact us for more information

contact@orionagriscience.co.nz

0800 674 6627